arXiv:0910.5358v1 [quant-ph] 28 Oct 2009

Nonclassicality of photon-added squeezed vacuum and its decoherence in thermal environment[*]

Li-yun Hu[1,2]†and Hong-yi Fan[2]
[1]College of Physics and Communication Electronics, Jiangxi Normal University, Nanchang 330022, China
[2]Department of Physics, Shanghai Jiao Tong University, Shanghai, 200030, China

October 28, 2009

Abstract

We study the nonclassicality of photon-added squeezed vacuum (PASV) and its decoherence in thermal environment in terms of the sub-Poissonian statistics and the negativity of Wigner function (WF). By converting the PASV to a squeezed Hermite polynomial excitation state, we derive a compact expression for the normalization factor of m-PASV, which is an m-order Legendre polynomial of squeezing parameter r. We also derive the explicit expression of WF of m-PASV and find the negative region of WF in phase space. We show that there is an upper bound value of r for this state to exhibit sub-Poissonian statistics increasing as m increases. Then we derive the explicit analytical expression of time evolution of WF of m-PASV in the thermal channel and discuss the loss of nonclassicality using the negativity of WF. The threshold value of decay time is presented for the single PASV.

PACS number(s): 03.67.-a, 03.65. Ud, 42. 50.Dv

1 Introduction

Nonclassical states play an important role in quantum computation and quantum imformation procession [1]. The nonclassicality of quantum states can be well-described by some nonclassical properties, such as sub-Poissonian photon ststistics [2], squeezing in one of the quadratures of the field [3], and negativity of Wigner function (WF) [4]. Especially, the partial negativity of WF implies the highly nonclassical properties of quantum states and is often used to describe the decoherence of quantum states. Many experimental schemes have been proposed to generate nonclassical states of optical field. Among them, subtracting photons form and/or adding photons to quantum states have been paid much attention because these fields exhibit an abundant of nonclassical properties and may give access to a complete engineering of quantum states and to fundamental quantum phenomena [5, 6, 7, 8, 10, 11, 12, 13]. For example, a single photon-subtraction squeezed vacuum (PSSV) has been experimentally prepared with a pulsed and continuous wave squeezed vacuum [11, 12, 13]. It is very similar to quantum superpositions of coherent states with small amplitudes [14, 15]. As another example, a single-photon addition was experimentally performed by Zavatta *et al* [6], which unveils the nonclassical features associated with the excitation of a classical coherent field by a single light quantum. For the single PSSV, its nonclassical properties and decoherence was investigated theoretically in two different decoherent channels (amplitude decay and phase damping) by Biswas and Agarwal [16]. They indicated that the WF losses its non-Gaussian nature and becomes Gaussian at long times in amplitude decay case.

---

[*]Project supported by the National Natural Science Foundation of China (Grant Nos 10775097 and 10874174) and the Research Foundation of the Education Department of Jiangxi Province.

[†]Corresponding author. *E-mail address*: hlyun2008@126.com (L-Y Hu).

On the other hand, the combination of the photon addition and subtraction has been successfully demonstrated in Ref.[8, 9]. In Ref.[8], photon addition and subtraction experimentally have been employed to probe quantum commutation rules by Parigi *et al*. In fact, they have implemented simple alternated sequences of photon creation (addition) and annihilation (subtraction) on a thermal field and observed the noncommutativity of the creation and annihilation operators. It is interesting to notice that subtracting or adding one photon from/to pure squeezed vacuums can generate the same output state, i.e., squeezed single-photon state [17]. However, the resulting states obtained by successive photon subtractions or additions are different from each other. For instance, successive two-photon additions $[a^{\dagger 2}]$ and successive two-photon subtractions $[a^2]$ will result in the same state produced by using subtraction-addition $(a^{\dagger}a)$ and addition-subtraction $(aa^{\dagger})$, (also see section 2 below) respectively. As far as we know, the nonclassicality and decoherence of arbitrary number photon-added squeezed vacuum states (PASV) in a dissipative channel has not been derived analytically in the literature before.

In this paper, we shall investigate the nonclassical properties and decoherence of single-mode PASV which is optically produced single-mode non-Gaussian states. This work is arranged as follows. In section 2, we introduce the single-mode PASV and discuss its nonclassicality in terms of sub-Poissonian statistics and the negativity of its Wigner function (WF). By converting the PASV to a squeezed Hermite polynomial excitation state, we derive a compact expression for the normalization factor of PASV, which is an m-order Legendre polynomial of the squeezing parameter r, where m is the number of added photons; and then derive the explicit analytical expression of WF for any photon-added number m and find the negative region of WF in phase space. We also show that there is an upper bound value of r for this state to exhibit sub-Poissonian statistics which increases as m increases. In section 3, we derive the explicit analytical expression of time evolution of WF of the arbitrary PASV in the thermal channel and discuss the loss of nonclassicality in reference of the negativity of WF. The threshold value of the decay time corresponding to the transition of the WF from partial negative to completely positive definite is obtained at the center of the phase space, which is independent of the squeezing parameter. We show that the WF for single PASV has always negative value for all parameters r if the decay time $\kappa t < \frac{1}{2}\ln\frac{2\bar{n}+2}{2\bar{n}+1}$ (see Eq.(40) below), where $\bar{n}$ denotes the average thermal photon number in the environment with dissipative coefficient κ. Conclusions are involved in the last section.

2 Single-mode Photon added squeezed vacuum state

Various photon states have been generated by the micromaser and WFs of some cavity fields can be measured by a scheme based on interaction between cavity fields and atoms [18, 19]. As described in Ref. [20] when an excited atom passes through a cavity field which is in a squeezed vacuum state then their interaction may produce photon addition (excitation) on the squeezed vacuum state—the excited squeezed vacuum state.

Therectically, the single-mode PASV can be obtained by repeatedly operating the photon creation operator $a^{\dagger}$ on a squeezed vacuum state $S(r)|0\rangle$, i.e.,

$$|r,m\rangle \equiv N_{r,m}a^{\dagger m}S(r)|0\rangle, \tag{1}$$

where $|0\rangle$ is single mode vacuum, and $N_{r,m}$ is the normalization constant to be determined, $a^{\dagger}$ is the Bose creation operator, and $S(r)$ is the single-mode squeezing operator $S(\lambda) = \exp[\frac{1}{2}\left(ra^{\dagger 2} - ra^2\right)]$ [21, 22] with r being the squeezing parameter.

2.1 Single-mode PASV as the squeezed Hermite polynomial excitation state

Under the transform of $S(\lambda)$ we see $S^{\dagger}(r)a^{\dagger}S(r) = a^{\dagger}\cosh r + a\sinh r$, thus we can reform Eq.(1) as

$$\begin{aligned}|r,m\rangle &= N_{r,m}S(r)S^{\dagger}(r)a^{\dagger m}S(r)|0\rangle \\ &= N_{r,m}S(r)\left(a^{\dagger}\cosh r + a\sinh r\right)^{m}|0\rangle.\end{aligned} \tag{2}$$

On the other hand, using the operator identity [23]

$$\left(a\mu+\nu a^{\dagger}\right)^{m}=\left(-i\sqrt{\frac{\mu\nu}{2}}\right)^{m}:H_{m}\left(i\sqrt{\frac{\mu}{2\nu}}a+i\sqrt{\frac{\nu}{2\mu}}a^{\dagger}\right):, \tag{3}$$

where $H_m(x)$ is m-order single variable Hermite polynomial whose definition is

$$H_{m}\left(x\right)=\sum_{l=0}^{[m/2]}\frac{\left(-1\right)^{l}m!}{l!(m-2l)!}\left(2x\right)^{m-2l},$$

we have

$$\begin{aligned}\left|r,m\right\rangle &=\frac{\left(-i\right)^{m}}{2^{m}}N_{r,m}\sinh^{m/2}2rS\left(r\right):H_{m}\left(i\sqrt{\frac{\tanh r}{2}}a+i\sqrt{\frac{\coth r}{2}}a^{\dagger}\right):\left|0\right\rangle \\ &=\frac{\left(-i\right)^{m}}{2^{m}}N_{r,m}\sinh^{m/2}2rS\left(r\right)H_{m}\left(i\sqrt{\frac{\coth r}{2}}a^{\dagger}\right)\left|0\right\rangle, \end{aligned} \tag{4}$$

which indicates that the PASV is equivalent to a squeezed Hermite-excited vacuum state.

Further using the generating function of $H_m(x)$,

$$H_{m}\left(x\right)=\left.\frac{\partial^{m}}{\partial t^{m}}\exp\left(2xt-t^{2}\right)\right|_{t=0}, \tag{5}$$

and Eq.(4), the normalization factor $N_{r,m}$ can be derived by

$$\begin{aligned}1&=\frac{N_{r,m}^{2}}{2^{2m}}\sinh^{m}2r\left\langle 0\right|H_{m}\left(-i\sqrt{\frac{\coth r}{2}}a\right)H_{m}\left(i\sqrt{\frac{\coth r}{2}}a^{\dagger}\right)\left|0\right\rangle \\ &=\frac{N_{r,m}^{2}}{2^{2m}}\sinh^{m}2r\frac{\partial^{2m}}{\partial\tau^{m}\partial t^{m}}e^{-t^{2}-\tau^{2}}\left.\left\langle 0\right|e^{-i\sqrt{2\coth r}a\tau}e^{i\sqrt{2\coth r}a^{\dagger}t}\left|0\right\rangle\right|_{\tau=t=0} \\ &=\frac{N_{r,m}^{2}}{2^{2m}}\sinh^{m}2r\frac{\partial^{2m}}{\partial\tau^{m}\partial t^{m}}\left.\exp\left(-t^{2}-\tau^{2}+2\tau t\coth r\right)\right|_{\tau=t=0}, \end{aligned} \tag{6}$$

where in the last step we have used the Baker-Hausdorff formula $e^{\mu a}e^{\nu a^{\dagger}}=e^{\nu a^{\dagger}}e^{\mu a}e^{\mu\nu}$. Using the newly found generating function of Legendre polynomial [24, 25] (see Appendix A),

$$\left.\frac{\partial^{2m}}{\partial t^{m}\partial\tau^{m}}\exp\left(-t^{2}-\tau^{2}+\frac{2x\tau t}{\sqrt{x^{2}-1}}\right)\right|_{t,\tau=0}=\frac{2^{m}m!}{\left(x^{2}-1\right)^{m/2}}P_{m}\left(x\right), \tag{7}$$

we can derive the compact expression for $N_{r,m}$, i.e.,

$$N_{r,m}^{-2}=m!\cosh^{m}rP_{m}\left(\cosh r\right), \tag{8}$$

which is just the result in Ref.[20] derived by the mathematical induction method. In particular, when $m=1$, ($H_1(x)=2x$) i.e., the single-photon added case, we see that $\left|r,1\right\rangle=S\left(r\right)a^{\dagger}\left|0\right\rangle$ is just a squeezed single-photon state; on the other hand, for the single-photon subtracted case [17, 26], the state is $aS\left(r\right)\left|0\right\rangle=S\left(r\right)S^{\dagger}\left(r\right)aS\left(r\right)\left|0\right\rangle=S\left(r\right)\left(a\cosh r+a^{\dagger}\sinh r\right)\left|0\right\rangle\rightarrow S\left(r\right)a^{\dagger}\left|0\right\rangle$, which indicates that adding a single-photon to the squeezed state has the same impact as annihilating a photon from the squeezed state. While for $m\geqslant 2$, the case is not true (also see Fig.1). For example, successive two-photon additions [$a^{\dagger 2}$] and successive two-photon subtractions [a^2] will result in the same state produced by using subtraction-addition ($a^{\dagger}a$) and addition-subtraction ($aa^{\dagger}$), respectively, i.e., $a^{\dagger 2}S\left(r\right)\left|0\right\rangle\rightarrow a^{\dagger}aS\left(r\right)\left|0\right\rangle, a^{2}S\left(r\right)\left|0\right\rangle\rightarrow aa^{\dagger}S\left(r\right)\left|0\right\rangle$. In Ref.[27], two PSSV is used to generate the squeezed superposition of coherent states with high fidelities and large amplitudes.

Combining Eqs.(1) and (8) we can conveniently calculate the average photon number $a^{\dagger}a$ in PASV,

$$\begin{aligned}\langle r,m|\, a^{\dagger}a\, |r,m\rangle &= \langle r,m|\, aa^{\dagger}\, |r,m\rangle - 1\\ &= \frac{N_{r,m}^{2}}{N_{r,m+1}^{2}} - 1\\ &= (m+1)\,\zeta\frac{P_{m+1}(\zeta)}{P_{m}(\zeta)} - 1,\end{aligned} \tag{9}$$

where we denote $\zeta = \cosh r$ for simplicity, and

$$\begin{aligned}\langle r,m|\, a^{\dagger 2}a^{2}\, |r,m\rangle &= \langle r,m|\left(a^{2}a^{\dagger 2} - 4aa^{\dagger} + 2\right)|r,m\rangle\\ &= \frac{N_{r,m}^{2}}{N_{r,m+2}^{2}} - 4\frac{N_{r,m}^{2}}{N_{r,m+1}^{2}} + 2\\ &= (m+1)\,\zeta\left\{(m+2)\,\zeta\frac{P_{m+2}(\zeta)}{P_{m}(\zeta)} - 4\frac{P_{m+1}(\zeta)}{P_{m}(\zeta)}\right\} + 2,\end{aligned} \tag{10}$$

thus the Mandel's $\mathcal{Q}$-parameter can be obtained by substituting Eqs.(9) and (10) into $\mathcal{Q} \equiv \frac{\langle a^{\dagger 2}a^{2}\rangle}{\langle a^{\dagger}a\rangle} - \left\langle a^{\dagger}a\right\rangle$. In particular, for single-photon-added case $m=1$, Eqs.(9) and (10) reduce to

$$\langle r,1|\, a^{\dagger}a\, |r,1\rangle = 3\cosh^{2} r - 2, \tag{11}$$

$$\langle r,1|\, a^{\dagger 2}a^{2}\, |r,1\rangle = \frac{3\left(3 + 2\tanh^{2} r\right)}{\left(\coth r - \tanh r\right)^{2}}, \tag{12}$$

thus the $\mathcal{Q}$-parameter with $m=1$ is given by

$$\mathcal{Q} = \frac{3\sinh^{2} 2r}{3\cosh 2r - 1} - 1. \tag{13}$$

From Eq.(13) we find that $\mathcal{Q}$ becomes negative for $m=1$ which is satisfied for the squeezing parameter $r \lesssim 0.46$ similar to the result of $\mathcal{Q}$ for single-photon subtracted case [16]. In order to see clearly the variation of $\mathcal{Q}$-parameter with r, we show the plots of $\mathcal{Q}$-parameter in Fig.1, from which one can clearly see that $\mathcal{Q}$-parameter becomes negetive $(m \neq 0)$ when r is less than a certain threshold value which increases as m increases; while for $m=0$, $\mathcal{Q}$ is always positive. This implies that the nonclassicality is enhanced by adding photon to squeezed state. We should emphasize that the WF has negative region for all r, and thus the PASV is nonclassical. In our following work, we pay attention to the (ideal) PASV in a thermal channel.

2.2 Wigner function of PASV

In order to discuss the decoherence properties of PASV in thermal environment, in this subsection we shall derive the analytical expression of Wigner function for PASVS. For single-mode case, the Wigner operator is defined as [28]

$$\Delta(q,p) = \frac{1}{2\pi}\int_{-\infty}^{\infty} du \left|q - \frac{u}{2}\right\rangle \left\langle q + \frac{u}{2}\right| e^{-ipu}, \tag{14}$$

where $|q\rangle$ is the coordinate representation, $Q\,|q\rangle = q\,|q\rangle$. Thus the Wigner function of PASVS $|r,m\rangle$ can be calculated by $W(q,p) = \langle r,m|\, \Delta(q,p)\, |r,m\rangle$. Using Eq.(4), we can see

$$W(q,p) = \frac{N_{r,m}^{2}}{2^{2m}}\sinh^{m} 2r\, \langle 0|\, H_{m}\left(-i\sqrt{\frac{\coth r}{2}}a\right) S^{\dagger}(r)\, \Delta(q,p)\, S(r)\, H_{m}\left(i\sqrt{\frac{\coth r}{2}}a^{\dagger}\right)|0\rangle. \tag{15}$$



Figure 1: (Color online) The Q-parameter as the function of squeezing parameter r for different $m = 0, 1, 2, 3, 4, 29, 30$.

On the other hand, noticing that the single-mode squeezing operator $S^{\dagger}(r)$ has its natural expression in coordinate representation [29], i.e., $S^{\dagger}(r) = e^{r/2}\int_{-\infty}^{\infty} dq \left|qe^{r}\right\rangle \left\langle q\right|,$ leading to $S^{\dagger}(r)\left|q\right\rangle = e^{-r/2}\left|qe^{-r}\right\rangle,$ it then follows that

$$\begin{aligned}
&S^{\dagger}(r)\,\Delta(q,p)\,S(r)\\
&=\frac{1}{2\pi}\int_{-\infty}^{\infty} d\left(ue^{-r}\right)\left|qe^{-r}-\frac{ue^{-r}}{2}\right\rangle\left\langle qe^{-r}+\frac{ue^{-r}}{2}\right| e^{-ipu}\\
&=\Delta\left(qe^{-r},pe^{r}\right).
\end{aligned}\tag{16}$$

Then substituting Eq.(16) into Eq.(15) and using Eq.(5) as well as the coherent state representation of Wigner operator,

$$\Delta(q,p)\rightarrow\Delta(\alpha,\alpha^{*})=\frac{e^{2|\alpha|^{2}}}{\pi}\int\frac{d^{2}z}{\pi}\left|z\right\rangle\left\langle -z\right| e^{-2(z\alpha^{*}-z^{*}\alpha)},\tag{17}$$

where $\alpha = (q+ip)/\sqrt{2}$ and $\left|z\right\rangle = \exp(za^{\dagger} - z^{*}a)\left|0\right\rangle$ is the coherent state [30, 31], we can put Eq.(15) into the following form,

$$\begin{aligned}
W(q,p) &= \frac{N_{r,m}^{2}}{2^{2m}}\sinh^{m}2r\left\langle 0\right| H_{m}\left(-i\sqrt{\frac{\coth r}{2}}a\right)\Delta\left(qe^{-r},pe^{r}\right)H_{m}\left(i\sqrt{\frac{\coth r}{2}}a^{\dagger}\right)\left|0\right\rangle\\
&=\frac{N_{r,m}^{2}}{2^{2m}}\sinh^{m}2r\frac{\partial^{2m}}{\partial t^{m}\partial\tau^{m}}e^{-t^{2}-\tau^{2}}\left\langle 0\right| e^{-i\sqrt{2\coth r}at}\Delta\left(qe^{-r},pe^{r}\right)e^{i\sqrt{2\coth r}a^{\dagger}\tau}\left|0\right\rangle\Big|_{\tau=t=0}\\
&=\frac{N_{r,m}^{2}e^{2|\beta|^{2}}}{2^{2m}\pi}\sinh^{m}2r\frac{\partial^{2m}}{\partial t^{m}\partial\tau^{m}}e^{-t^{2}-\tau^{2}}\int\frac{d^{2}z}{\pi}e^{-|z|^{2}-\left(i\sqrt{2\coth r}t+2\beta^{*}\right)z+\left(2\beta-i\sqrt{2\coth r}\tau\right)z^{*}}\Big|_{\tau=t=0}\\
&=\frac{N_{r,m}^{2}e^{-2|\beta|^{2}}}{2^{2m}\pi}\sinh^{m}2r\,\frac{\partial^{2m}}{\partial t^{m}\partial\tau^{m}}e^{-t^{2}+2\bar{\beta}t-\tau^{2}+2\bar{\beta}^{*}\tau-2t\tau\coth r}\Big|_{\tau=t=0},
\end{aligned}\tag{18}$$

where

$$\bar{\beta}=-i\beta\sqrt{2\coth r},\beta=\left(qe^{-r}+ipe^{r}\right)/\sqrt{2}=\alpha\cosh r-\alpha^{*}\sinh r,\tag{19}$$

and in the last step, we have used the integration formula

$$\int\frac{d^{2}z}{\pi}e^{\zeta|z|^{2}+\xi z+\eta z^{*}}=-\frac{1}{\zeta}e^{-\frac{\xi\eta}{\zeta}},\mathrm{Re}\left(\zeta\right)<0.\tag{20}$$

In order to further simplify Eq.(18), expanding the exponential item $e^{-2t\tau\coth r}$ as sum series and using Eq.(5) we have

$$\begin{aligned}W(q,p) &= \frac{N_m^2 e^{-2|\beta|^2}}{2^{2m}\pi}\sinh^m 2r\sum_{l=0}^{\infty}\frac{(-2\coth r)^l}{2^{2l}l!}\frac{\partial^{2l}}{\partial(\bar{\beta})^l\,\partial(\bar{\beta}^*)^l}\left.\frac{\partial^{2m}}{\partial t^m\partial\tau^m}e^{-t^2+2\bar{\beta}t-\tau^2+2\bar{\beta}^*\tau}\right|_{\tau=t=0}\\ &= \frac{N_m^2 e^{-2|\beta|^2}}{2^{2m}\pi}\sinh^m 2r\sum_{l=0}^{\infty}\frac{(-2\coth r)^l}{2^{2l}l!}\frac{\partial^{2l}}{\partial(\bar{\beta})^l\,\partial(\bar{\beta}^*)^l}H_m(\bar{\beta})H_m(\bar{\beta}^*). \qquad (21)\end{aligned}$$

Noticing the recurrence relations of $H_m(x)$,

$$\frac{\mathrm{d}}{\mathrm{d}x^l}H_m(x) = \frac{2^l m!}{(m-l)!}H_{m-l}(x), \qquad (22)$$

then the Wigner function of $|r, m\rangle$ is given by

$$\begin{aligned}W(q,p) &= \frac{N_m^2 e^{-2|\beta|^2}}{2^{2m}\pi}\sinh^m 2r\sum_{l=0}^{m}\frac{(m!)^2(-2\coth r)^l}{l!\,[(m-l)!]^2}\left|H_{m-l}(\bar{\beta})\right|^2\\ &= \frac{1}{\pi}\frac{e^{-2|\beta|^2}\sinh^m r}{2^m P_m(\cosh r)}\sum_{l=0}^{m}\frac{m!\,(-2\coth r)^l}{l!\,[(m-l)!]^2}\left|H_{m-l}(\bar{\beta})\right|^2, \qquad (23)\end{aligned}$$

where β and $\bar{\beta}$ are shown in Eq.(19). Eq.(23) seems a new result (not reported in the literature before), related to single-variable Hermite polynomials. In particular, when the photon-added number $m = 0, 1$ and noticing that $P_0(\cosh r) = 1$ and $P_1(\cosh r) = \cosh r$, Eq.(23) reduce to, respectively,

$$W_{m=0}(q,p) = \frac{1}{\pi}e^{-(q^2e^{-2r}+p^2e^{2r})}, \qquad (24)$$

$$W_{m=1}(q,p) = \frac{1}{\pi}\left\{2(q^2e^{-2r}+p^2e^{2r})-1\right\}e^{-(q^2e^{-2r}+p^2e^{2r})}. \qquad (25)$$

Eq.(24) is just the WF of squeezed vacuum state, a Gaussian in phase space; while Eq.(25) corresponds to a non-Gaussian WF in phase space due to the presence of non-Gaussian item $2(q^2e^{-2r} + p^2e^{2r}) - 1$. It is clear that the function $W_{m=1}(q,p)$ becomes negative in phase space, when $2(q^2e^{-2r} + p^2e^{2r}) < 1$.

Using Eq.(23) we show the plots of WF in the phase space in Figs.2 for different squeezing parameters r and photon-added numbers m. One can see clearly that there is some negative region of the WF in the phase space which implies the nonclassicality of this state. In addition, the squeezing effect in one of the quadratures is clear in the plots (see Figs.2a and Figs.2b), which is another evidence of the nonclassicality of this state. The WF has its minimum value for $m = 1, 3$ at the center of phase space $(q = p = 0)$ (see Fig.2(a) and (d)). The case is not true for $m = 2$ (see Fig2.c).

3 Decoherence of PASV in thermal environment

3.1 Model of Decoherence

In this section, we consider how this single-mode state (1) evolves at the presence of thermal environment. In thermal channel, the evolution of the density matrix for the m-PASV can be described by [32]

$$\frac{d\rho}{dt} = \kappa(\bar{n}+1)\left(2a\rho a^\dagger - a^\dagger a\rho - \rho a^\dagger a\right) + \kappa\bar{n}\left(2a^\dagger\rho a - aa^\dagger\rho - \rho aa^\dagger\right), \qquad (26)$$

where κ represents the dissipative coefficient and $\bar{n}$ denotes the average thermal photon number of the environment. When $\bar{n} = 0$, Eq.(26) reduces to the master equation describing the photon-loss channel. The corresponding time evolution density operator can be obtained as

$$\rho_{r,m}(t) = e^{\kappa t+\Gamma_0}\sum_{k,l=0}^{\infty}M_{k,l}\rho_0 M_{k,l}^\dagger, \qquad (27)$$



Figure 2: (Color online) The Wigner functions of PASV for (a) $m=1, r=0.3$,(b) $m=1, r=0.8$,(c) $m=2, r=0.3$ and (d) $m=3, r=0.3$.

where $\rho_0 = |r,m\rangle\langle r,m|$ is the initial density matrix; $M_{k,l}$ and $M_{k,l}^{\dagger}$ are Hermite conjugated operators (Kraus operator) with each other,

$$M_{k,l} = e^{(\kappa t+\Gamma_0)/2}\sqrt{\frac{\Gamma_-^k \Gamma_+^l e^{-2l\Gamma_0}}{k!l!}}\, e^{\Gamma_0 a^{\dagger} a} a^{\dagger l} a^k, \tag{28}$$

and Γ_+, Γ_- and Γ_0 are determined by

$$T = 1 - e^{-2\kappa t},\ \Gamma_+ = \frac{\bar{n}T}{\bar{n}T+1},\ \Gamma_- = \frac{(\bar{n}+1)\,T}{\bar{n}T+1},\ \Gamma_0 = \ln\frac{e^{-\kappa t}}{\bar{n}T+1}. \tag{29}$$

It is not difficult to prove the $M_{k,l}$ obeys the normalization condition $\sum_{k,l=0}^{\infty} M_{k,l}^{\dagger} M_{k,l} = 1$ by using the technique of integration within an ordered products of operators [33, 34].

3.2 Wigner function of the PASV in a thermal channel

The evolution formula of WF of the PASV can be derived as follows [35]

$$W(\zeta,\zeta^*,t) = \frac{2}{(2\bar{n}+1)\,T}\int\frac{d^2\alpha}{\pi} W(\alpha,\alpha^*,0)\, e^{-2\frac{\left|\zeta-\alpha e^{-\kappa t}\right|^2}{(2\bar{n}+1)T}}, \tag{30}$$

where $W(\alpha,\alpha^*,0)$ is the WF of the initial state. Eq.(30) is just the evolution formula of WF in thermal channel. Thus the WF at any time can be obtained by performing the integration when the initial WF is known.

In a similar way to deriving Eq.(23), substituting Eq.(23) into Eq.(30) and using the generating function of single-variable Hermite polynomials (5) and Eq.(20), we finally obtain (see appendix B)

$$\begin{aligned} W(\zeta,\zeta^*,t) &= \frac{2\sinh^m r}{\pi\sqrt{C}\,(2\bar{n}+1)\,T}\frac{m!e^{-\frac{2|\zeta|^2}{(2\bar{n}+1)T}}}{2^m P_m(\cosh r)} e^{\frac{A}{C}|B|^2+\frac{\sinh 2r}{C}\left(B^{*2}+B^2\right)} \\ &\times\sum_{l=0}^{m}\sum_{k=0}^{m-l}\frac{(-2\coth r)^l\, G^{m-l-k}F^k}{l!k!\left[(m-l-k)!\right]^2}\left|H_{m-l-k}(E/\sqrt{G})\right|^2. \end{aligned} \tag{31}$$

where we have set

$$A=\frac{2e^{-2\kappa t}}{\left(2\bar{n}+1\right)T}+2\cosh 2r, \tag{32}$$

$$B=\frac{2e^{-\kappa t}\zeta}{\left(2\bar{n}+1\right)T},\ C=A^{2}-4\sinh^{2}2r, \tag{33}$$

$$D=\sqrt{2\coth r}i\left(B^{*}\sinh r-B\cosh r\right), \tag{34}$$

$$E=\frac{1}{C}\left(AD-2D^{*}\sinh 2r\right), \tag{35}$$

$$F=\frac{8}{C}\left(A\cosh 2r\coth r-4\cosh^{2}r\sinh 2r\right), \tag{36}$$

$$G=1-\frac{16}{C}\frac{e^{-2\kappa t}}{\left(2\bar{n}+1\right)T}\cosh^{2}r. \tag{37}$$

Eq.(31) is just the analytical expression of WF for the PASV in thermal channel. It is obvious that the WF loss its Gaussian property due to the presence of single-variable Hermite polynomials. In particular, at the initial time ($t=0$), noting $\sqrt{C}\left(2\bar{n}+1\right)T\rightarrow 2$, $G\rightarrow 1$, $F/G\rightarrow 0$, and $E/\sqrt{G}\rightarrow\bar{\beta}=-i\sqrt{2\coth r}\left(\zeta\cosh r-\zeta^{*}\sinh r\right)$, as well as $\frac{A}{C}\left|B\right|^{2}-\frac{2|\zeta|^{2}}{(2\bar{n}+1)T}\rightarrow -2\left|\zeta\right|^{2}\cosh 2r$, $\frac{\sinh 2r}{C}\left(B^{*2}+B^{2}\right)\rightarrow\left(\zeta^{2}+\zeta^{*2}\right)\sinh 2r$, Eq.(31) just does reduce to Eq.(23), i.e., the WF of the PASV. On the other hand, when $\kappa t\rightarrow\infty$, noticing that $T\rightarrow 1, B\rightarrow 0, C\rightarrow 4, D\rightarrow 0,\ E/\sqrt{G}\rightarrow 0, G\rightarrow 1,\ F\rightarrow 4\coth r$,as well as $H_{m}\left(0\right)=\left(-1\right)^{j}\frac{m!}{j!}\delta_{m,2j}$, then Eq.(31) becomes $\frac{1}{\pi(2\bar{n}+1)}e^{-\frac{2|\zeta|^{2}}{2\bar{n}+1}}$, which is independent of photon-addition number m and corresponds to the WF of thermal state with mean thermal photon number $\bar{n}$. This indicates that the system state reduces to a thermal state after an enough long time interaction with the environment.

In addition, for the case of $m=0$, single-mode squeezed vacuum, Eq.(31) just becomes ($H_{0}(x)=1$)

$$W_{m=0}\left(\zeta,\zeta^{*},t\right)=\mathfrak{N}e^{-\mathfrak{D}|\zeta|^{2}+\mathfrak{E}(\zeta^{*2}+\zeta^{2})}, \tag{38}$$

where $\mathfrak{N}=\frac{2}{\pi\sqrt{C}(2\bar{n}+1)T}$ is the normalization factor, $\mathfrak{D}=\frac{2}{(2\bar{n}+1)T}-\frac{A\mathfrak{E}}{\sinh 2r}$, $\mathfrak{E}=\frac{4e^{-2\kappa t}\sinh 2r}{[(2\bar{n}+1)T]^{2}C}$, Eq.(38) denotes a Gaussian distribution function— the WF of single-mode squeezed vacuum in the thermal channel; while for $m=1$, single–photon added case, its WF in the thermal channel is given by ($H_{1}(x)=2x$)

$$W_{m=1}\left(\zeta,\zeta^{*},t\right)=\frac{4\left|E\right|^{2}+F-2\coth r}{\pi\sqrt{C}\left(2\bar{n}+1\right)T\coth r}e^{\frac{A}{C}|B|^{2}-\frac{2|\zeta|^{2}}{(2\bar{n}+1)T}+\frac{\sinh 2r}{C}\left(B^{*2}+B^{2}\right)}. \tag{39}$$

In Fig.3, the WFs of the PASV with $m=1$ are depicted in phase space with $r=0.3$ and $\bar{n}=1$ for several different κt. It is easy to see that the negative region of WF gradually diminishes as the time κt increases. Actually, from Eq.(33) one can see that $C>0$, so at the center of the phase space ($\alpha=\alpha^{*}=0$), when $F<2\coth r$ leading to the following condition:

$$\kappa t<\kappa t_{c}\equiv\frac{1}{2}\ln\frac{2\bar{n}+2}{2\bar{n}+1}, \tag{40}$$

which is independent of the squeezing parameter r, there always exist negative region for WF in phase space and the WF of PASV is always positive in the whole phase space when κt exceeds the threshold value κt_{c}.

In Figs. 4 and 5, we plot the variation of WF in phase space for different $\bar{n}$ and r, respectively. It is found that the partial negativity of WF decreases gradually as $\bar{n}$ (or r) increases for a given time. The squeezing effect in one of the quadrature is shown in Fig.5. In addition, for the case of large squeezing value r, the single-photon added squeezed state becomes similar to a Schodinger cat state (see Fig.6). The WF becomes Gaussian with the time evolution. In principle, by using the explicit expression of WF in Eq.(31), we can draw the WF distribution for any photon-added case in phase



Figure 3: (Color online) The Wigner functions of single-photon-added squeezed vacuum states in phase space for $r = 0.3, \bar{n} = 1$ at (a) $\kappa t = 0.05$, (b) $\kappa t = 0.1$, (c) $\kappa t = 0.2$ and (d) $\kappa t = 0.5$.

space. For instance, for $m = 2$, there are two negative regions of the WF, which differs from the case of single PASV (see Fig.7). The absolute value of the negative minimum of the WF decreases as κt increases, which leads to the complete absence of partial negative region.

4 Conclusions

The nonclassical properties and decoherence of single-mode PASV in a thermal environment have been investigated. A compact expression for the normalization factor of PASV is derived by converting the PASV to a squeezed Hermite polynomial excitation state. It is shown that the normalization factor is just an m-order Legendre polynomial of the squeezing parameter r. We also derived the explicit analytical expression of WF for any photon-added number m and found the negative region of WF in phase space. We also show that there is an upper bound value of r for this state to exhibit sub-Poissonian statistics which increases as m increases. Then we considered the effects of decoherence to the nonclassicality of PASV when interacting with thermal environment. For arbitrary number PASV, we derived the explicit analytical expression of time evolution of WF and presented the loss of nonclassicality in reference of the negativity of WF. The threshold value of the decay time corresponding to the transition of the WF from partial negative to completely positive definite is obtained. We find that the WF has always negative value for all parameters r if the decay time $\kappa t < \frac{1}{2} \ln \frac{2\bar{n}+2}{2\bar{n}+1}$ for single PASV.

ACKNOWLEDGEMENTS: Work supported by the National Natural Science Foundation of China (Grant Nos 10775097 and 10874174) and the Research Foundation of the Education Department of Jiangxi Province.



Figure 4: (Color online) The Wigner functions of single-photon-added squeezed vacuum states in phase space for $r = 0.3$ and $\kappa t = 0.05$ with (a) $\bar{n} = 0, (b)$ $\bar{n} = 1, (c)$ $\bar{n} = 2, (d)$ $\bar{n} = 5$.



Figure 5: (Color online) The Wigner functions of single-photon-subtracted squeezed vacuum states in phase space for $\bar{n} = 0.1$ and $\kappa t = 0.05$ with (a) $r = 0.03, (b)$ $r = 0.5, (c)$ $r = 0.8, (d)$ $r = 1.5.??$



Figure 6: (Color online) Wigner function of single photon-added squeezed vacuum states in phase space for $r = 0.8, \bar{n} = 0$: (a) $\kappa t = 0.1, (b)$ $\kappa t = 0.2, (c)$ $\kappa t = 0.3$, and (d) $\kappa t = 0.7$.



Figure 7: (Color online) The Wigner functions of photon-added squeezed vacuum states in phase space for $r = 0.7, \bar{n} = 1, m = 2$: (a) $\kappa t = 0.1$, (b) $\kappa t = 0.2$.

Appendix A: Derivation of Eq.(7)

Recalling that the newly found expression of Legendre polynomial [24, 25]

$$x^m \sum_{l=0}^{[m/2]} \frac{m!}{2^{2l} (l!)^2 (m-2l)!} \left(1 - \frac{1}{x^2}\right)^l = P_m(x), \tag{A1}$$

which is equivalent to the well-known Legendre polynomial's expression [36]

$$P_m(x) = \sum_{l=0}^{[m/2]} (-1)^l \frac{(2m-2l)!}{2^m l! (m-l)! (m-2l)!} x^{m-2l}, \tag{A2}$$

though it is different in form from Eq.(A1), they actually are equal. For instance, what we list in the following equations (the left) from Eq.(A1) are equal to what we list from (A2) (the right)

$$\begin{aligned} &m = 0,\ P_0(x) = 1; \\ &m = 1,\ P_1(x) = x; \\ &m = 2,\ P_2(x) = x^2 \left[1 + \frac{1}{2}\left(1 - \frac{1}{x^2}\right)\right] = \frac{3}{2}x^2 - \frac{1}{2}; \\ &m = 3,\ P_3(x) = x^3 \left[1 + \frac{3}{2}\left(1 - \frac{1}{x^2}\right)\right] = \frac{5}{2}x^3 - \frac{3}{2}x; \end{aligned} \tag{A3}$$

and

$$\begin{aligned} &m = 4,\ P_4(x) = x^4 \left[1 + 3\left(1 - \frac{1}{x^2}\right) + \frac{3}{8}\left(1 - \frac{1}{x^2}\right)^2\right] = \frac{1}{8}\left(35x^4 - 30x^2 + 3\right); \\ &m = 5,\ P_5(x) = x^5 \left[1 + 5\left(1 - \frac{1}{x^2}\right) + \frac{15}{8}\left(1 - \frac{1}{x^2}\right)^2\right] = \frac{1}{8}\left(63x^5 - 70x^3 + 15x\right); \end{aligned} \tag{A4}$$

We emphasize that the new form in Eq.(A1) cannot be directly obtained by some series summation rearrangement technique from the orginal definition (A2).

On the other hand, noting the following relation

$$\begin{aligned} &\frac{\partial^{2m}}{\partial t^m \partial \tau^m} \exp\left(-t^2 - \tau^2 + 2x\tau t\right)\Big|_{t,\tau=0} \\ &= \sum_{n,l,k=0}^{\infty} \frac{(-)^{n+l}}{n! l! k!} (2x)^k \left.\frac{\partial^{2m}}{\partial t^m \partial \tau^m} \tau^{2n+k} t^{2l+k}\right|_{t,\tau=0} \\ &= 2^m m! \sum_{n=0}^{[m/2]} \frac{m!}{2^{2n} (n!)^2 (m-2n)!} x^{m-2n}, \end{aligned} \tag{A5}$$

and then comparing Eq.(A5) with Eq.(A1) we can obtain Eq.(7).

APPENDIX B: Derivation of Wigner function Eq.(31) of PASV

In this appendix, we present the details for deriving the Wigner function Eq.(31). Substituting Eq.(23) into Eq.(30) and noticing Eq.(19) as well as the generating function of single-variable Hermite polynomials (5), we have

$$\begin{aligned} W\left(\zeta,\zeta^{*},t\right) &= \bar{N}\sum_{l=0}^{m}\frac{\left(-2\coth r\right)^{l}}{l!\left[(m-l)!\right]^{2}}\int\frac{d^{2}\alpha}{\pi}e^{-2\frac{\left|\zeta-\alpha e^{-\kappa t}\right|^{2}}{(2\bar{n}+1)T}-2\left|\alpha\cosh r-\alpha^{*}\sinh r\right|^{2}} \\ &\quad\times\left|H_{m-l}(-i\left(\alpha\cosh r-\alpha^{*}\sinh r\right)\sqrt{2\coth r})\right|^{2} \\ &= \bar{N}e^{-\frac{2\left|\zeta\right|^{2}}{(2\bar{n}+1)T}}\sum_{l=0}^{m}\frac{\left(-2\coth r\right)^{l}}{l!\left[(m-l)!\right]^{2}}\frac{\partial^{2m-2l}}{\partial\upsilon^{m-l}\partial\tau^{m-l}}e^{-\upsilon^{2}-\tau^{2}} \\ &\quad\times\int\frac{d^{2}\alpha}{\pi}\exp\left\{-A\left|\alpha\right|^{2}+B_{1}\alpha+B_{2}\alpha^{*}+\left(\alpha^{2}+\alpha^{*2}\right)\sinh 2r\right\}\Big|_{\tau=\upsilon=0}, \end{aligned} \tag{B1}$$

where A and B are given by Eqs.(34)-(35),

$$\bar{N}=\frac{2\sinh^{m}r}{\pi\left(2\bar{n}+1\right)T}\frac{m!}{2^{m}P_{m}\left(\cosh r\right)}, \tag{B2}$$

and

$$B_{1}=B^{*}-i2\left(\upsilon\cosh r+\tau\sinh r\right)\sqrt{2\coth r}, \tag{B3}$$

$$B_{2}=B+i2\left(\upsilon\sinh r+\tau\cosh r\right)\sqrt{2\coth r}. \tag{B4}$$

Further using the following integral identity [37]:

$$\begin{aligned} &\int\frac{d^{2}z}{\pi}\exp\left(\zeta\left|z\right|^{2}+\xi z+\eta z^{*}+fz^{2}+gz^{*2}\right) \\ &=\frac{1}{\sqrt{\zeta^{2}-4fg}}\exp\left[\frac{-\zeta\xi\eta+\xi^{2}g+\eta^{2}f}{\zeta^{2}-4fg}\right], \end{aligned} \tag{B5}$$

whose convergent condition is Re$(\zeta\pm f\pm g)<0$, and Re$\left(\frac{\zeta^{2}-4fg}{\zeta\pm f\pm g}\right)<0$, we can put Eq.(B1) into the following form:

$$\begin{aligned} W\left(\zeta,\zeta^{*},t\right) &= \frac{\bar{N}}{\sqrt{C}}e^{-\frac{2\left|\zeta\right|^{2}}{(2\bar{n}+1)T}}\sum_{l=0}^{m}\frac{\left(-2\coth r\right)^{l}}{l!\left[(m-l)!\right]^{2}}\frac{\partial^{2m-2l}}{\partial\upsilon^{m-l}\partial\tau^{m-l}} \\ &\quad\times\exp\left\{-\upsilon^{2}-\tau^{2}+\frac{A}{C}B_{1}B_{2}+\frac{1}{C}\left(B_{1}^{2}+B_{2}^{2}\right)\sinh 2r\right\}_{\tau=\upsilon=0}, \end{aligned} \tag{B6}$$

where C and D are given by (34)-(35), and

$$B_{1}B_{2}=B^{*}B+2\left(\upsilon D+\tau D^{*}\right)+8\left(\tau^{2}+\upsilon^{2}\right)\cosh^{2}r+8\tau\upsilon\cosh 2r\coth r, \tag{B7}$$

$$B_{1}^{2}+B_{2}^{2}=B^{*2}+B^{2}-32\tau\upsilon\cosh^{2}r-8\left(\upsilon^{2}+\tau^{2}\right)\cosh 2r\coth r-4\tau D-4\upsilon D^{*}, \tag{B8}$$

then substituting Eqs.(B7)-(B8) into Eq.(B6) yields

$$\begin{aligned} W\left(\zeta,\zeta^{*},t\right) &= \frac{\bar{N}}{\sqrt{C}}e^{-\frac{2\left|\zeta\right|^{2}}{(2\bar{n}+1)T}}e^{\frac{A}{C}\left|B\right|^{2}+\frac{\sinh 2r}{C}\left(B^{*2}+B^{2}\right)}\sum_{l=0}^{m}\frac{\left(-2\coth r\right)^{l}}{l!\left[(m-l)!\right]^{2}} \\ &\quad\times\frac{\partial^{2m-2l}}{\partial\upsilon^{m-l}\partial\tau^{m-l}}\exp\left\{-G\left(\upsilon^{2}+\tau^{2}\right)+2\upsilon E+2\tau E^{*}+\tau\upsilon F\right\}_{\tau=\upsilon=0}, \end{aligned} \tag{B9}$$

where E, F, and G are given by Eqs.(35)-(37). In order to further simplify Eq.(B9), expanding the exponential item $e^{\tau\upsilon F}$ as sum series and using Eqs.(5) and the formula

$$\left.\frac{\partial^m}{\partial\upsilon^m}e^{-G\upsilon^2+2\upsilon E}\right|_{\upsilon=0}=G^{m/2}H_m\left(E/\sqrt{G}\right), \tag{B10}$$

we see

$$\begin{aligned}W\left(\zeta,\zeta^*,t\right)&=\frac{\bar{N}}{\sqrt{C}}e^{-\frac{2|\zeta|^2}{(2\bar{n}+1)T}+\frac{A}{C}|B|^2+\frac{\sinh 2r}{C}\left(B^{*2}+B^2\right)}\sum_{l=0}^{m}\frac{\left(-2\coth r\right)^l}{l!\left[(m-l)!\right]^2}\\&\times\sum_{k=0}^{\infty}\frac{F^k}{k!}\frac{\partial^{2k}}{\partial\left(2E\right)^k\partial\left(2E^*\right)^k}\frac{\partial^{2m-2l}}{\partial\upsilon^{m-l}\partial\tau^{m-l}}\left.e^{-G\left(\upsilon^2+\tau^2\right)+2\upsilon E+2\tau E^*}\right|_{\tau=\upsilon=0}\\&=\frac{\bar{N}}{\sqrt{C}}e^{-\frac{2|\zeta|^2}{(2\bar{n}+1)T}+\frac{A}{C}|B|^2+\frac{\sinh 2r}{C}\left(B^{*2}+B^2\right)}\sum_{l=0}^{m}\frac{\left(-2\coth r\right)^l G^{m-l}}{l!\left[(m-l)!\right]^2}\\&\times\sum_{k=0}^{\infty}\frac{F^k}{k!}\frac{\partial^{2k}}{\partial\left(2E\right)^k\partial\left(2E^*\right)^k}\left|H_{m-l}\left(E/\sqrt{G}\right)\right|^2.\end{aligned} \tag{B11}$$

Then using Eq.(22) yields Eq.(31).

References

[1] D. Bouwmeester, A. Ekert and A. Zeilinger, *The Physics of Quantum Information* (Springer-Verlag, Berlin, 2000).

[2] R. Short and L. Mandel, Phys. Rev. Lett. **51**, 384 (1983).

[3] V. V. Dodonov, J. Opt. B: Quantum Semiclassical Opt. **4**, R1 (2002).

[4] M. Hillery, R. F. O'Connell, M. O. Scully, and E. P. Wigner, Phys. Rep. **106**, 121 (1984).

[5] J. Wenger, R. Tualle-Brouri, and P. Grangier, Phys. Rev. Lett. 92, 153601 (2004).

[6] A. Zavatta, S. Viciani, and M. Bellini, Science, **306**, 660 (2004).

[7] A. Zavatta, S. Viciani, and M. Bellini, Phys. Rev. A **72**, 023820 (2005).

[8] V. Parigi, A. Zavatta, M. S. Kim, and M. Bellini, Science, **317**, 1890 (2007).

[9] R. W. Boyd, K. W. Chan, and M. N. O'Sullivan, Science, **317**, 1874 (2007).

[10] A. Zavatta, S. Viciani, and M. Bellini, Phys. Rev. A **75**, 052106 (2007).

[11] J. S. Neergaard-Nielsen, B. Melholt Nielsen, C. Hettich, K. Mølmer, and E. S. Polzik, Phys. Rev. Lett. 97, 083604 (2006).

[12] A. Ourjoumtsev, R. Tualle-Brouri, J. Laurat, Ph. Grangier, Science **312**, 83 (2006).

[13] K. Wakui, H. Takahashi, A. Furusawa, and M. Sasaki, Opt. Express **15**, 3568 (2007).

[14] M. Dakna, T. Anhut, T. Opatrny, L. Knoll, and D.-G. Welsch, Phys. Rev. A **55**, 3184 (1997).

[15] S. Glancy and H. M. de Vasconcelos, J. Opt. Soc. Am. B **25**, 712 (2008).

[16] A. Biswas and G. S. Agarwal, Phys. Rev. A **75**, 032104 (2007).

[17] H. Jeong, A. P. Lund, and T. C. Ralph, Phys. Rev. A **72**, 013801 (2005).

[18] B. T. H. Vracoe, S. Brattke, M. Weidinger, H. Walther, Nature **403**, 743 (2000);
S. Brattke, B.T.H. Vracoe, H. Walther, Phys. Rev. Lett. **86**, 3534 (2001).

[19] Z. M. Zhang, Chin. Phys. Lett. 20 (2003) 227;
Z. M. Zhang, Chin. Phys. Lett. 21 (2004) 5.

[20] Z. X. Zhang, H. Y. Fan, Phys. Lett. A 174 (1993) 206.

[21] D. F. Walls andG J Milburn, *Quantum Optics* (Springer-Verlag, Berlin, 1994).

[22] M. O. Scully and M. S. Zubairy, *Quantum Optics* (Cambridge: Cambidge University Press, 1997).

[23] H.Y. Fan and Vander J. Linde, J. Phys. A **24**, 2529 (1989).

[24] L. Y. Hu and H. Y. Fan, J. Opt. Soc. Am. B, **25**, 1955 (2008).

[25] H. Y. Fan, X. G. Meng and J. S. Wang, Commun. Theor. Phys. **46,** 845 (2006).

[26] M. S. Kim, J. Phys. B: At. Mol. Opt. Phys. **41**, 133001 (2008).

[27] P. Marek, H. Jeong, and M. S. Kim, Phys. Rev. A 78, 063811 (2008).

[28] H.-Y. Fan, H. R. Zaidi, Phys. Lett. A 124, 303 (1987).

[29] H.-Y. Fan, Representation and Transformation Theory in Quantum Mechanics, Shanghai Scientific & Technical, Shanghai Press, 1997.

[30] R. Glauber, Phys. Rev. **130,** 2529 (1963).

[31] R. Glauber, Phys. Rev. **131,** 2766 (1963).

[32] C. Gardiner and P. Zoller, *Quantum Noise* (Springer, Berlin, 2000).

[33] H.-Y. Fan, H.-L. Lu and Y. Fan, Ann. Phys. **321,** 480 (2006).

[34] H.-Y. Fan and L. Y. Hu, Mod. Phys. Lett. B, 22, 2435 (2008).

[35] L. Y. Hu and H.-Y. Fan, Opt. Commun. **282**, 4379 (2009).

[36] I. S. Gradshteyn and L. M. Ryzhik, *Tables of Integration Series and Products* (Academic Press, New York, 1980).

[37] R. R. Puri, *Mathematical Methods of Quantum Optics* (Springer-Verlag, Berlin, 2001), Appendix A.